

Cefokel 50 mg/ml

suspension for injection 

Composition

Ceftiofur (as hydrochloride) 50 mg –
excipients up to 1 ml.

Target species

Cattle and pigs.

Indications

Infections associated with bacteria sensitive to ceftiofur:

• In cattle:

For the treatment of bacterial respiratory disease associated with *Mannheimia haemolytica* (previously *Pasteurella haemolytica*), *Pasteurella multocida* and *Histophilus somni* (previously *Haemophilus somnus*).

For the treatment of acute interdigital necrobacillosis (panaritium, foot rot), associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus* (*Porphyromonas asaccharolytica*).

For treatment of the bacterial component of acute post-partum (puerperal) metritis within 10 days after calving associated with *Escherichia coli*, *Arcanobacterium pyogenes* and *Fusobacterium necrophorum*, sensitive to ceftiofur. The indication is restricted to cases where treatment with another antimicrobial has failed.

• In pigs:

For the treatment of bacterial respiratory disease associated with *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *Streptococcus suis*.

Dosage and administration

Cattle:

Respiratory disease: 1 mg ceftiofur /kg b.w./day for 3 to 5 days by subcutaneous injection, i.e.

1 ml/50 kg b.w. at each injection.

Acute interdigital necrobacillosis:

1 mg/kg b.w./day for 3 days by subcutaneous injection, i.e. 1 ml/50 kg b.w. at each injection.

Acute post-partum metritis within 10 days after calving:

1 mg/kg b.w./day for 5 consecutive days by subcutaneous injection, i.e. 1 ml/50 kg b.w. at each injection.

Before use, shake the bottle vigorously for at least 30 seconds until the product appears adequately resuspended.

The recommended maximum volume to be administered at a single injection site is 4 ml in pigs and 6 ml in cattle. Subsequent injections must be given at different sites.

The vial cannot be broached more than 66 times.

In case of acute post-partum metritis, additional supportive therapy might be required in some cases.

Pigs:

3 mg ceftiofur /kg b.w./day for 3 days via intramuscular route, i.e. 1 ml/16 kg b.w. at each injection.

Pack size

Glass vials of 100 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

