

# Combikel 40 L.A.

suspension for injection 

## Composition

Procaine benzylpenicillin 120 000 I.U. –  
Benzathine benzylpenicillin 80 000 I.U. –  
Dihydrostreptomycin sulphate eq. 200 mg base –  
Excipients up to 1 ml.

## Target species

Cattle, horses, pigs, goats, sheep, dogs, cats.

## Indications

Prevention and treatment of infections caused by penicillin G and/or DHS susceptible microorganisms such as respiratory tract infections (shipping fever, bronchopneumonias, upper respiratory tract infections), uro-genital tract infections (e.g. pyelonephritis by *Corynebacterium renale*, endometritis), infections of skin and wounds, streptococci infections in horses (strangles, *Str. zooepidemicus* infections in foals), erysipelas in pigs, panaritium, necrobacillosis in calves and piglets, anaerobe infections (clostridial infections, tetanus, anthrax), polyarthritis (navel ill), mastitis, septicaemia, bacterial infections secondary to viral ones. The drug also can be used in postoperative prevention of infections.

## Dosage and administration

By intramuscular or subcutaneous administration. Shake before use.

### Dosage:

- **Cattle:** 5 - 10 ml of Combikel 40 L.A. per 100 kg b.w.
- **Horses, pigs, goats, sheep:** 5 ml of Combikel 40 L.A. per 50 kg b.w.
- **Dogs, cats:** 0.5 ml of Combikel 40 L.A. per 5 kg b.w.

If necessary, treatment should be repeated after 3 days.

## Pack size

100 ml, 250 ml.

### Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

