

# Combikel 40

suspension for injection 

## Composition

Procaine benzylpenicillin (lecithin coated)  
200 mg (eq. 200 000 I.U.)  
Dihydrostreptomycin sulphate eq. 200 mg  
Dihydrostreptomycin  
Excipients up to 1 ml.

## Target species

Horses, cattle, sheep, goats, dogs, cats.

## Indications

Infections caused by penicillin G and/or DHS susceptible microorganisms such as respiratory tract infections (shipping fever, pasteurellosis, bronchopneumonia, pleuropneumonia, rhinitis, laryngitis), uro-genital tract infections (e.g. pyelonephritis, nephritis, endometritis), infections of skin and wounds, peritonitis, strangles in horses, mastitis, meningitis (e.g. by streptococci), erysipelas in pigs, otitis in dogs and cats, panaritium, clostridial infections, tetanus, anthrax, leptospirosis, actinomycosis in cattle, actinobacillosis in horses, polyarthritis (navel ill) and septicaemia.

The drug also can be used in postoperative prevention of infections (e.g. caesarean section).

## Dosage and administration

By intramuscular or subcutaneous route.  
Shake before use.

Dosage: 200 000 I.U. of penicillin G and 200 mg dihydrostreptomycin (base) or 1 ml of the suspension per 10 - 20 kg of body weight once daily.

## Pack size

Vials of 50 and 100 ml.

### Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

