

Dexakel 02

Injectable solution 

Composition

Dexamethasone sodium phosphate eq. 2 mg
dexamethasone
Excipients up to 1 ml.

Target species

Cattle, horses, sheep, goats, calves, dogs, cats.

Indications

- Metabolic disorders such as: ketosis in cattle and puerperal toxemia in ewes and sows.
- Non-infectious inflammatory processes, especially acute musculoskeletal inflammations such as: arthritis, peri-arthritis, tendovaginitis, bursitis, luxations, myositis, osselets and sprains.
- As an aid in acute infectious diseases (e.g. acute mastitis, metritis, pneumonia, polyarthritis in calves) in combination with suitable anti-infectious therapy.
- Allergic conditions, e.g. asthma, allergic skin affections (eczema, urticaria, pruritis, ...), snake-bites.
- Stress- and shock conditions.
- Induction of parturition in ruminants during the last stage of pregnancy (delivery within 2-4 days).

Dosage and administration

By intravenous, intramuscular or subcutaneous (dogs, cats) injection:

- **Cats, dogs:**
(0.1 - 0.2 mg/kg b.w.) 0.25 - 0.5 ml/5 kg
- **Cattle, horses:**
(2.5 - 5 mg/100 kg b.w.) 5 - 10 ml/400 kg
- **Sheep, goats, calves:**
(2 - 4 mg/50 kg) 1 - 2 ml/50 kg

If necessary repeat with 2 days intervals in dogs and cats and 3-4 days intervals in other animals.

- For induction of parturition in cows (given within 1 week before expected calving date):
(5 mg/100 kg b.w.) 2.5 ml/100 kg.
- Puerperal toxemia and induction of parturition in sheep and goats:
(2.5 mg/10 kg b.w.) 6 ml/50 kg.
- Shock-therapy: by slow I.V. injection of high doses (10 x), eventually to be repeated after 8-12 h.

By intra-articular injection: according to the extension of joint and animal weight: 0.25 to 4 ml.

Pack size

50 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

