

Diminakel PLUS

granulate for injection 

Composition

1 g of powder contains:
 Diminazene diacetate 445 mg
 Vitamin B12 0.64 mg
 Vitamin B6 2.54 mg
 Antipyrine 551.82 mg.

Target species

Cattle, horses, camels, dogs.

Indications

For the treatment and prevention of trypanosomiasis (nagana, surra, dourine).

For the treatment of babesiosis.

For the treatment of theileriosis caused by *Theileria annulata*.

Dosage and administration

Dissolve 2.36 g of powder in 12.5 ml of distilled water to obtain a ready-to-use 7 % solution (70 mg/ml).

Subcutaneous or intramuscular administration, the latter is recommended in horses.

a) Treatment of trypanosomiasis:

- **horses, cattle:** 3.5 mg of diminazene diacetate/kg b.w. (1 ml/20 kg b.w.).
- **camels:** 1.25 mg of diminazene diacetate/kg b.w. (1 ml/55 kg b.w.).

b) Treatment of babesiosis:

- **horses, cattle, dogs:** 3.5 mg of diminazene diacetate/kg b.w. (1 ml/20 kg b.w.).

c) Treatment of theileriosis:

- **cattle:** 5 mg of diminazene diacetate/kg b.w. (1 ml/14 kg b.w.).

Pack size

10 sachets of 2.36 g or 23.6 g.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

