

Diminakel

granules for injection 

Composition

1 g of powder contains: Diminazene diacetate 445 mg – Antipyrine 555 mg.

Target species

Cattle, sheep, camels, horses, dogs.

Indications

For the treatment of babesiosis and theileriosis in animals.

For the sterilisation of babesia infections in ruminants and horses to be exported or moved to disease free areas.

For the treatment of canine malignant icterus.

For the treatment and prevention of trypanosomiasis in animals.

Dosage and administration

One sachet of 2.36 g powder is sufficient for 300 kg b.w. at a dose of 3.5 mg diminazene diacetate/kg b.w.

Dissolve 1 sachet of 2.36 g powder in 12.5 ml (or 1 sachet of 23.6 g in 125 ml) of sterile water for injection to obtain a ready-to-use solution of 7% (= about 15 ml).

Inject the solution by subcutaneous or intramuscular routes. Use preferably the I.M. route in horses.

a) Treatment of babesiosis:

- **Cattle, sheep, dogs:**
3.5 mg diminazene diacetate/kg b.w.
or 1 ml/20 kg b.w.
- **Horses:**
5 mg diminazene diacetate/kg b.w.
or ± 10 ml/140 kg b.w.
on 2 consecutive days.

b) Treatment of theileriosis:

- **Cattle:**
5 - 10 mg diminazene diacetate/kg b.w.
or 10 ml/70 - 140 kg b.w.,
for 2 consecutive days
or 3 - 4 injections on alternate days.

c) Treatment of trypanosomiasis:

- **Cattle, horses and dogs:**
T. congolense, T. vivax:
3.5 mg diminazene diacetate/kg b.w.
or 1 ml/20 kg b.w.
T. brucei, T. evansi, T. equiperdum:
7 mg diminazene diacetate/kg b.w.
or 1 ml/10 kg b.w.
- **Camels:**
1.75 mg diminazene diacetate/kg b.w.
or 1 ml/40 kg b.w.

Pack size

10 sachets of 2.36 g or 23.6 g.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

