

Dovikel

injectable and oral solution 

Composition

Nitroxinil 250 mg
Excipients up to 1 ml.

Target species

Cattle, sheep, goats, rabbits, camels, pheasants.

Indications

- **Cattle:**
 - adult *Fasciola hepatica* (partially active against larval stages)
 - adult *Fasciola gigantica*
 - gastro-intestinal nematodes: *Haemonchus spp.*, *Bunostomum spp.*, *Oesophagostomum spp.*
- **Sheep and goats:**
 - adult *Fasciola hepatica* (partially active against larval stages)
 - adult *Fasciola gigantica*
 - Gastro-intestinal nematodes: *Haemonchus spp.*, ivermectin- and benzimidazol resistant *Haemonchus contortus*
 - Larvae of the nasal bot fly (*Oestrus*)
- **Rabbits:** *Fasciola hepatica*, *Dicrocoelium dentriticum*
- **Pheasants:** *Syngamus trachea*.

Dosage and administration

Fascioliasis:

Subcutaneous injection in cattle, sheep, goats and camels of 10 mg/kg or 1 ml per 25 kg b.w.
In acute phase (larval stages):
13 mg/kg or 1.3 ml per 25 kg b.w.

Nematodes:

Subcutaneous injection in cattle, sheep and goats of 12.5 mg/kg b.w. or 1.25 ml per 25 kg b.w.

Oestrus ovis-Larvae:

Subcutaneous injection of 15 - 20 mg/kg b.w. or 1.5 - 2.0 ml per 25 kg b.w.

Rabbits: subcutaneous injection of 10 mg/kg b.w. or 0.04 ml/kg b.w. (1 ml of the solution can be diluted with 9 ml of water for injection: then inject 0.4 ml/kg b.w.)

Pheasants: oral administration using 0.5 ml per litre drinking water.

Pack size

50 ml, 250 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

