

Florfenikel 300 mg/ml

injectable solution 

Composition

Florfenicol 300 mg – Excipients up to 1 ml.

Target species

Cattle and pigs.

Indications

Diseases caused by florfenicol susceptible bacteria.

Pigs:

Treatment of acute outbreaks of swine respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol.

Cattle:

Treatment of respiratory tract infections due to strains of *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* susceptible to florfenicol.

Dosage and mode of administration

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing. The vial cannot be broached more than 25 times.

Pigs:

IM injection of 15 mg/kg body weight (1 ml/20 kg) into the neck muscle twice 48 hours apart. The volume administered per injection site should not exceed 3 ml. Subsequent injections must be given at different sites.

Cattle:

IM injection of 20 mg/kg body weight (1ml/15 kg) into the neck muscle twice 48 hours apart. The volume administered per injection site should not exceed 10 ml. Subsequent injections must be given at different sites.

Pack size

Vials of 100 and 250 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

