

Gentakel 05

Injectable solution 

Composition

Gentamicin sulphate 50 000 I.U. (= 50 mg base)
Excipients up to 1 ml.

Target species

Cattle, pig, dog.

Indications

For treatment of infectious diseases, caused by Gram-positive and Gram-negative bacteria susceptible to gentamicin.

Dosage and mode of administration

Cattle and pigs should be administered intramuscularly. Dogs should be administered intramuscularly or subcutaneously.

Cattle, pigs:

2-3 mg gentamicin / kg body weight, i.e.,
1-1.5 ml of preparation / 25 kg bodyweight,
administered solely by intramuscular use.
Particularly during the first day of treatment, it is
advised to administer this dose twice with a
12 hours interval. The duration of treatment
should not exceed 3 days.
Up to 1 ml of formulation can be given to
pigs at one injection site!

Dogs:

4 mg gentamicin / kg body weight, i.e.,
1 ml of preparation / 12.5 kg body weight.
This dose is given twice daily with an interval of
12 hours on the first day of treatment and once
daily thereafter. The average duration of treat-
ment is 4-7 days, depending on the severity of
the disease, with a maximum of 14 days.

Pack size

Vials of 50 ml and 100 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

