



Composition

Oxytetracycline hydrochloride eq. 100 mg base Excipients up to 1 ml.

Target species

Cattle, sheep, goats, pigs

Indications

Infections caused by oxytetracycline sensitive germs, such as:

Cattle:

- treatment and prevention of respiratory infections
- treatment of foot rot, otitis media

Sheep, goats:

- respiratory infections
- treatment of foot rot, omphalitis

Pigs:

- treatment and prophylaxis of respiratory infections (enzootic abortion, atrophic rhinitis, infections caused by Pasteurella and Haemophilus spp.).
- treatment and prophylaxis of Mastitis-Metritis-Agalactic Syndrome (MMA) in sows.

Dosage and administration

By deep I.M. injection

Calves:

1 ml/5 kg b.w. every 24-48 hrs.

Cattle:

0,5-1 ml/10 kg b.w. every 24 hrs.

Pigs, goats, sheep:

1-2 ml/10 kg b.w. every 24 hrs.

Repeat the dosage as necessary for 2-3 injections at 24-48 hours intervals. No more than 20 ml should be administered at one injection site.

Pack size

Vials of 50 ml and 100 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.









