

Penikel 300

suspension for injection 

Composition

Procaine benzylpenicillin (lecithin coated)
300 mg (eq. 300 000 I.U.) Excipients up to 1 ml.

Target species

Cattle, pigs.

Indications

Treatment of infections caused by procaine benzylpenicillin susceptible bacteria. One must consider the diffusion of the antibiotic at the infection site in sufficient concentration.

Cattle: respiratory infections caused by *Pasteurella spp.*; Mastitis caused by *Streptococcus dysgalactiae*, *A. pyogenes*, penicillin G-sensitive *Staphylococcus aureus*; urinary or genital tract infections caused by *Arcanobacterium pyogenes*; umbilical cord infections caused by *Arcanobacterium pyogenes*.

Pigs: respiratory infections caused by *Pasteurella spp.*, *Actinobacillus pleuropneumoniae*; urinary or genital tract infections (incl. MMA syndrome) caused by Beta-hemolytic streptococci; Swine Erysipelas; central nervous system or musculoskeletal system infections caused by *Arcanobacterium pyogenes* or *Streptococcus suis*.

Dosage and administration

Shake bottle before use.

Intramuscular administration: 21 000 I.U./kg b.w. or 0.7 ml/10 kg b.w., once a day during maximum 5 days in cattle and max. 3 days in pigs

Pack size:

Vials of 100 ml and 250 ml.

Disclaimer

Please be aware that product names, product specifications and availability can differ between countries. Not all products and indications are licensed in every country. For more detailed information in this regard, please contact us.

