

# AMOXYCILLIN 150 LA

**COMPOSITION:**

Amoxicillin 150 LA (Amoxicillin 150 mg/ml) contains per ml: Amoxicillin trihydrate eq. 150 mg amoxicillin.

**TARGET SPECIES:**

Cattle, pigs, sheep, dogs, cats.

**INDICATIONS:**

For the treatment of infections caused by Gram-positive and Gram-negative bacteria that are susceptible to amoxicillin.

**CONTRA-INDICATIONS:**

Do not use in animals with a history of hypersensitivity towards penicillins and cephalosporins.  
Do not use in small rodents (Guinea pigs, hamsters, gerbils, rabbits) since amoxicillin may cause fatal colitis in these species.

**DOSAGE:**

7.5 to 15 mg of amoxicillin/kg/b.w./day or 0.5 to 1 ml Amoxicillin 150 LA/10 kg b.w./day. The dose may be repeated if necessary. A dose interval of 36-48 hours is recommended.

**METHOD AND ROUTES OF ADMINISTRATION:**

- Dogs, cats: by intramuscular or subcutaneous administration.
- Cattle, pigs, sheep: by intramuscular route only.

**UNDESIRABLE EFFECTS:**

Allergic reactions (skin; life-threatening anaphylaxia) may occur in hypersensitized animals.

**SPECIAL PRECAUTIONS FOR USE, OTHER WARNINGS:**

Shake before use. Do not use by intravenous route. Use always a dry syringe.  
Use during pregnancy and lactation is not contra-indicated.  
Amoxicillin 150 LA should not be mixed with other drugs in the same syringe.

**PACK SIZE:**

100 ml.

