

ANCOPEN P

COMPOSITION:

One injector contains:
Kanamycin sulphate eq. 100 mg base
Procaine benzylpenicillin 300 000 I.U.
Prednisolone 20 mg
Excipients up to 10 g.

PHARMACOLOGICAL PROPERTIES:

Ancopen P contains 3 active components, namely the synergistic association of the antibiotics procaine penicillin G and kanamycin sulphate which show antimicrobial activity against most important udder pathogens (*Streptococcus agalactiae*, *Str. dysgalactiae*, *Str. uberis*, *Staphylococcus aureus*, *Actinomyces pyogenes*, *E. coli*) and the glucocorticoid prednisolone. Addition of the anti-inflammatory short acting prednisolone makes this preparation particularly of value in treatment of acute mastitis, often accompanied by painful tissue reactions.

INDICATIONS:

For treatment of acute mastitis in cows, sheep and goats.

CONTRA-INDICATIONS:

Do not use in animals known to be hypersensitive to penicillins.

DOSAGE AND ADMINISTRATION:

Prior to treatment, milk out the udder thoroughly. Clean and disinfect teats. Thereafter insert the cannula into the teat canal and inject the ointment. The contents should be massaged upwards. If necessary, repeat the treatment with intervals of 12-24 hours.

Use one injector per infected cow quarter.

In ewes, use 2 injectors per infected mammary gland.

PACK SIZE:

Box with 20 injectors.